

<https://doi.org/10.33380/3034-3925-2026-3-2-65-66>



## Ресурсный потенциал *Onosma simplicissima* L. (Boraginaceae), произрастающей в Новосибирской области

Д. С. Круглов, Е. Д. Олешко, В. В. Величко, Д. Л. Прокушева

Гербариум. 2026;3(1):33–40. <https://doi.org/10.33380/3034-3925-2026-3-1-63>. Статья опубликована: 16.01.2026.

На странице 34 в исходную версию статьи редакция внесла изменение в Abstract в разделы "Introduction" и "Materials and methods".

### Вместо:

**Introduction.** Wild medicinal plants represent an important source of biologically active compounds widely used in pharmaceutical practice. Species of the genus *Onosma* L. (Boraginaceae) are characterized by high ecological plasticity and significant pharmacological potential associated with the accumulation of phenolic compounds. *Onosma simplicissima* L., widely distributed in the steppe and forest-steppe landscapes of Western Siberia, is considered a promising source of flavonoids and oxycinnamic acids. However, an adequate assessment of its resource potential requires consideration of both biomass reserves and environmentally driven variability of the phytochemical profile.

**Aim.** To perform a comprehensive assessment of the resource potential of *Onosma simplicissima* L. within the Novosibirsk Region and to determine the influence of habitat conditions on the content of phenolic compounds in herb raw material.

**Materials and methods.** The study was conducted on the aerial parts of *O. simplicissima* collected at the flowering stage from nine locations in the Novosibirsk Region. Resource characteristics were evaluated using the model plant method with calculation of biological and exploitable reserves as well as potential annual harvest volumes. The total content of oxycinnamic acids and flavonoids was determined by spectrophotometry and expressed as rosmarinic acid and rutin equivalents, respectively. Statistical analysis was performed using Student's t-test at a confidence level of  $P = 95\%$ .

### Исправлено на:

**Introduction.** Wild medicinal plants represent an important source of biologically active compounds widely used in pharmaceutical practice. Species of the genus *Onosma* L. (Boraginaceae) are characterized by high ecological plasticity and significant pharmacological potential associated with the accumulation of phenolic compounds. *Onosma simplicissima* L., widely distributed in the steppe and forest-steppe landscapes of Western Siberia, is considered a promising source of flavonoids and oxycinnamic acids. However, an adequate assessment of its resource potential requires consideration of both biomass reserves and environmentally driven variability of the phytochemical profile.

**Aim.** To perform a comprehensive assessment of the resource potential of *Onosma simplicissima* L. within the Novosibirsk Region and to determine the influence of habitat conditions on the content of phenolic compounds in herb raw material.

**Materials and methods.** The study was conducted on the aerial parts of *O. simplicissima* collected at the flowering stage from nine locations in the Novosibirsk Region. Resource characteristics were evaluated using the model plant method with calculation of biological and exploitable reserves as well as potential annual harvest volumes. The total content of oxycinnamic acids and flavonoids was determined by spectrophotometry and expressed as rosmarinic acid and rutin equivalents, respectively. Statistical analysis was performed using Student's t-test at a confidence level of  $P = 95\%$ .

Внесение исправление не повлияло на содержание статьи.

Обновлена онлайн-версия статьи на сайте журнала.

## Resource Potential of *Onosma simplicissima* L. (Boraginaceae) Growing in the Novosibirsk Region

Dmitriy S. Kruglov, Egor D. Oleshko, Victoria V. Velichko, Daria L. Prokusheva

Herbarium. 2026;3(1):33–40. (In Russ.) <https://doi.org/10.33380/3034-3925-2026-3-1-63>. Published: 16.01.2026.

On page 34 of the original version of the article, the editors made a change to the Abstract in the "Introduction" and "Materials and methods" sections.

**Instead:**

**Introduction.** Wild medicinal plants represent an important source of biologically active compounds widely used in pharmaceutical practice. Species of the genus *Onosma* L. (*Boraginaceae*) are characterized by high ecological plasticity and significant pharmacological potential associated with the accumulation of phenolic compounds. *Onosma simplicissima* L., widely distributed in the steppe and forest-steppe landscapes of Western Siberia, is considered a promising source of flavonoids and oxycinnamic acids. However, an adequate assessment of its resource potential requires consideration of both biomass reserves and environmentally driven variability of the phytochemical profile.

**Aim.** To perform a comprehensive assessment of the resource potential of *Onosma simplicissima* L. within the Novosibirsk Region and to determine the influence of habitat conditions on the content of phenolic compounds in herb raw material.

**Materials and methods.** The study was conducted on the aerial parts of *O. simplicissima* collected at the flowering stage from nine locations in the Novosibirsk Region. Resource characteristics were evaluated using the model plant method with calculation of biological and exploitable reserves as well as potential annual harvest volumes. The total content of oxycinnamic acids and flavonoids was determined by spectrophotometry and expressed as rosmarinic acid and rutin equivalents, respectively. Statistical analysis was performed using Student's t-test at a confidence level of  $P = 95\%$ .

**Corrected to:**

**Introduction.** Wild medicinal plants represent an important source of biologically active compounds widely used in pharmaceutical practice. Species of the genus *Onosma* L. (*Boraginaceae*) are characterized by high ecological plasticity and significant pharmacological potential associated with the accumulation of phenolic compounds. *Onosma simplicissima* L., widely distributed in the steppe and forest-steppe landscapes of Western Siberia, is considered a promising source of flavonoids and oxycinnamic acids. However, an adequate assessment of its resource potential requires consideration of both biomass reserves and environmentally driven variability of the phytochemical profile.

**Aim.** To perform a comprehensive assessment of the resource potential of *Onosma simplicissima* L. within the Novosibirsk Region and to determine the influence of habitat conditions on the content of phenolic compounds in herb raw material.

**Materials and methods.** The study was conducted on the aerial parts of *O. simplicissima* collected at the flowering stage from nine locations in the Novosibirsk Region. Resource characteristics were evaluated using the model plant method with calculation of biological and exploitable reserves as well as potential annual harvest volumes. The total content of oxycinnamic acids and flavonoids was determined by spectrophotometry and expressed as rosmarinic acid and rutin equivalents, respectively. Statistical analysis was performed using Student's t-test at a confidence level of  $P = 95\%$ .

The correction did not affect the content of the article.

The online version of the article on the journal's website has been updated.